

EVEN AND ODD KAUFFMAN BRACKET IDEALS FOR GENUS-1 TANGLES

SUSAN M. ABERNATHY AND PATRICK M. GILMER

ABSTRACT. We adapt a basis of Habiro's for the even Kauffman bracket skein module of $S^1 \times D^2$ to define bases for the even and odd skein modules of $S^1 \times D^2$ relative to two points. We discuss genus-1 tangle embedding, and define an even and odd version of the previously defined Kauffman bracket ideal for genus-1 tangles. These even and odd Kauffman bracket ideals are obstructions to tangle embeddings. Using our even and odd bases for the relative skein modules, we show how to compute a finite list of generators for the even and odd Kauffman bracket ideals of a genus-1 tangle. We do this explicitly for three genus-1 tangles. We relate these ideals to determinants of closures of genus-1 tangles.

1. INTRODUCTION

Let $M \subseteq S^3$ be a compact, oriented 3-manifold with boundary. Then an $(M, 2n)$ -tangle is 1-manifold with $2n$ boundary components properly embedded in M . We refer to $(S^1 \times D^2, 2)$ -tangles where $S^1 \times D^2$ is a unknotted solid torus in S^3 as genus-1 tangles.

An $(M, 2n)$ -tangle \mathcal{T} embeds in a link $L \subseteq S^3$ if there exists a complementary 1-manifold \mathcal{T}' with $2n$ boundary components in $S^3 - \text{Int}(M)$ such that upon gluing \mathcal{T}' to \mathcal{T} along their boundaries, we obtain a link isotopic to L . Such a link is called a closure of \mathcal{T} . We refer to \mathcal{T}' as the complementary 1-manifold of the closure. Note that if \mathcal{T} is a genus-1 tangle, then \mathcal{T}' is also a genus-1 tangle. The focus of this paper is genus-1 tangle embedding.

In [A, A3], the first author defined the notion of even and odd closures for any genus-1 tangle \mathcal{G} with respect to a longitude l on the boundary of the solid torus which misses the boundary points of \mathcal{T} . If we choose l to be the longitude pictured in Figure 1 and assume that

Date: July 25, 2015.

2010 Mathematics Subject Classification. 57M25.

Key words and phrases. tangles, tangle embedding, determinants, Kauffman bracket skein module.

The first author was supported as a research assistant by NSF-DMS-1311911.

The second author was partially supported by NSF-DMS-1311911.

the boundary points of \mathcal{T} are in the complement of l , then we may think of even and odd closures intuitively as follows. Even (respectively, odd) closures are those whose complementary 1-manifold passes through the hole of the solid torus containing \mathcal{G} an even (respectively, odd) number of times. For the remainder of this paper, when we discuss even and odd closures, we mean even and odd with respect to the longitude l .

In [A2, A3], the first author defined the Kauffman bracket ideal of an $(M, 2n)$ -tangle \mathcal{T} to be the ideal $I_{\mathcal{T}}$ of $\mathbb{Z}[A, A^{-1}]$ generated by the reduced Kauffman bracket polynomials of all closures of \mathcal{T} which gave an obstruction to embedding. In the case $(M, 2n) = (B^3, 4)$, this ideal was first studied by Przytycki, Silver and Williams [PSW].


The first author outlined a method for computing this ideal in the case of genus-1 tangles using skein theory techniques. In this paper, we define an even and odd version of the Kauffman bracket ideal for genus-1 tangles. The even Kauffman bracket ideal of a genus-1 tangle \mathcal{G} is the ideal $I_{\mathcal{G}}^{\text{even}}$ generated by the reduced Kauffman bracket polynomials of all even closures of \mathcal{G} . The odd Kauffman bracket ideal $I_{\mathcal{G}}^{\text{odd}}$ is defined similarly. If an ideal is equal to $\mathbb{Z}[A, A^{-1}]$, we refer to that ideal as trivial.

The following proposition is an immediate consequence of these definitions.

Proposition 1.1. *Let \mathcal{G} be a genus-1 tangle. If $I_{\mathcal{G}}^{\text{even}}$ (respectively, $I_{\mathcal{G}}^{\text{odd}}$) is non-trivial, then the unknot is not an even (respectively, odd) closure of \mathcal{G} . More generally, if L is an even (respectively, odd) closure of \mathcal{G} , then the reduced Kauffman bracket polynomial of L must lie in $I_{\mathcal{G}}^{\text{even}}$ (respectively, $I_{\mathcal{G}}^{\text{odd}}$).*

We also note that if the ordinary Kauffman bracket ideal is non-trivial, then both the even and odd Kauffman bracket ideals must be non-trivial. However, the converse is not true. In §6 we give an example of a genus-1 tangle (see Figure 6) which has non-trivial even and odd Kauffman bracket ideals but trivial ordinary Kauffman bracket ideal. We also show that Krebs' tangle [K], pictured in Figure 1, has trivial even ideal and non-trivial odd ideal, and give another example (in Figure 4) which has non-trivial even ideal and trivial odd ideal. In [A2, A3], we showed that the ordinary Kauffman bracket ideal of Krebs' tangle \mathcal{A} is trivial.

In §5, we outline a method for computing a finite list of generators for the even and odd Kauffman bracket ideals, following the same basic procedure as in [A2, A3]. We use the graph basis defined in [A2, A3] as well as bases for the even and odd Kauffman bracket skein modules of $S^1 \times D^2$ relative to two points which we define in §3. These even

FIGURE 1. Krebs' tangle \mathcal{A} and a longitude l .

and odd bases are adapted from a definition of Habiro's [Hab]. In the §7, we relate the even and odd ideals to the determinants of even and odd closures, connecting the results of this paper to those of [A].

2. KAUFFMAN BRACKET SKEIN MODULES

The Kauffman bracket polynomial of a framed link D , denoted by $\langle D \rangle$, is an element of $\mathbb{Z}[A, A^{-1}]$ given by the following three relations, where $\delta = -A^2 - A^{-2}$:

- (i) $\langle \text{crossing} \rangle = A \langle \text{positive crossing} \rangle + A^{-1} \langle \text{negative crossing} \rangle$
- (ii) $\langle D' \amalg \bigcirc \rangle = \delta \langle D' \rangle$.
- (iii) $\langle \rangle = 1$.

We let $\langle D \rangle'$ denote the reduced Kauffman bracket polynomial of D ; that is, where $\langle D \rangle' = \langle D \rangle / \delta$.

The Kauffman bracket skein module of a 3-manifold M is the $\mathbb{Z}[A, A^{-1}]$ -module $K(M)$ generated by isotopy classes of framed links in M modulo the Kauffman bracket relations above.

Of particular concern to us is the relative Kauffman bracket skein module. Let M be a compact oriented 3-manifold with boundary and a set of m specified marked framed points on ∂M . Then the Kauffman bracket skein module of M relative to the m marked points is the $\mathbb{Z}[A, A^{-1}]$ -module $K(M, m)$ generated by isotopy classes of framed 1-manifolds with boundary the marked framed points modulo the above Kauffman bracket relations. We can view any genus-1 tangle (equipped with the blackboard framing) as a skein element in $K(S^1 \times D^2, 2)$.

As in [A2, A3], we generalize the Hopf pairing on $K(S^1 \times D^2)$ defined in [BHMV] to obtain the relative Hopf pairing $\langle , \rangle : K(S^1 \times D^2, 2) \times K(S^1 \times D^2, 2) \rightarrow K(S^3) = \mathbb{Z}[A, A^{-1}]$. Given a and b in $K(S^1 \times D^2, 2)$,

we let

$$\langle a, b \rangle = \left\langle \begin{array}{c} \text{diagram of two circles with arcs } a \text{ and } b \end{array} \right\rangle$$

where a and b lie in regular neighborhoods of the trivalent graphs.

If a genus-1 tangle \mathcal{G} embeds in a link $L \subseteq S^3$, then we can describe this tangle embedding using the relative Hopf pairing. We have that $\langle L \rangle = \langle \mathcal{G}, \mathcal{G}' \rangle$ for some $\mathcal{G}' \in K(S^1 \times D^2, 2)$.

3. EVEN AND ODD RELATIVE SKEIN MODULES

As in [BHMV], we let z denote a standard banded core of $S^1 \times D^2$ and the element this core represents in $K(S^1 \times D^2)$. As described in [Hab], one can obtain a \mathbb{Z}_2 -graded algebra structure on the Kauffman bracket skein module $K(S^1 \times D^2)$ by letting $K^{\text{even}}(S^1 \times D^2)$ be the subalgebra of $K(S^1 \times D^2)$ generated by z^2 and $K^{\text{odd}}(S^1 \times D^2)$ be $zK^{\text{even}}(S^1 \times D^2)$. Then, one has that $K(S^1 \times D^2) = K^{\text{even}}(S^1 \times D^2) \oplus K^{\text{odd}}(S^1 \times D^2)$. This is because the Kauffman skein relations respect \mathbb{Z}_2 -homology classes [GH, p.105].

Suppose now that $S^1 \times D^2$ is equipped with two marked framed points in $\partial(S^1 \times D^2)$ and an essential curve l in $\partial(S^1 \times D^2)$ missing the marked points and which bounds a disk \mathfrak{D} in $S^1 \times D^2$. Let u be a framed 1-manifold in $S^1 \times D^2$ with the two given marked points as boundary. Then we say that u is even (respectively, odd), if u intersects \mathfrak{D} an even (respectively, odd) number of times. Let $K^{\text{even}}(S^1 \times D^2, 2)$ and $K^{\text{odd}}(S^1 \times D^2, 2)$ be the submodules of $K(S^1 \times D^2, 2)$ generated by all even and odd 1-manifolds, respectively. Then, we have that $K(S^1 \times D^2, 2) = K^{\text{even}}(S^1 \times D^2, 2) \oplus K^{\text{odd}}(S^1 \times D^2, 2)$. Note that if L is an even closure of a genus-1 tangle \mathcal{G} , then the Kauffman bracket polynomial of L can be written as $\langle L \rangle = \langle \mathcal{G}, \mathcal{G}' \rangle$ where $\mathcal{G} \in K(S^1 \times D^2, 2)$ and $\mathcal{G}' \in K^{\text{even}}(S^1 \times D^2, 2)$. Here l is a “longitude” for the first copy of $S^1 \times D^2$ and a “meridian” for the second copy of $S^1 \times D^2$. The analogous statement is true for odd closures.

We let $\phi_k = -A^{2k+2} - A^{-2k-2}$ (in [BHMV] and elsewhere this is denoted λ_i). The authors define a basis of $K(S^1 \times D^2)$ in [BHMV] as follows. Let $Q_n = \prod_{i=0}^{n-1} (z - \phi_i)$ for $n \geq 0$. Habiro modifies this definition




FIGURE 2. An even element of $K(S^1 \times D^2, 2)$, where \mathfrak{D} is the shaded disk

in [Hab] to obtain a basis for the even submodule $K^{\text{even}}(S^1 \times D^2)$ by defining $S_n = \prod_{i=0}^{n-1} (z^2 - \phi_i^2)$ for $n \geq 0$. Note that $S_n = Q_n \prod_{i=0}^{n-1} (z + \phi_i)$.

We adapt Habiro's basis to obtain bases for the $K^{\text{even}}(S^1 \times D^2, 2)$ and $K^{\text{odd}}(S^1 \times D^2, 2)$. We refer to them as the even basis and odd basis, respectively, and define them as follows (using the same \mathfrak{D} as pictured in Figure 2). The even basis consists of the following elements, where $n \geq 0$:

$$x_n^{\text{even}} = \text{diagram with } S_n \text{ label} \quad \text{and} \quad y_n^{\text{even}} = \text{diagram with } zS_n \text{ label}.$$

Similarly, the odd basis consists of the following elements, where $n \geq 0$:

$$x_n^{\text{odd}} = \text{diagram with } zS_n \text{ label} \quad \text{and} \quad y_n^{\text{odd}} = \text{diagram with } S_n \text{ label}.$$

That these are bases follows ultimately from the basis for $K(S^1 \times D^2, 2)$ consisting of framed links described by isotopy classes of diagrams without crossings and without contractible loops in $S^1 \times D^2$. One also uses the fact that there is a triangular unimodular change of basis matrix over $\mathbb{Z}[A, A^{-1}]$ relating the bases $\{S_n\}$ and $\{z^{2m}\}$ for $K^{\text{even}}(S^1 \times D^2)$.

4. GRAPH BASIS OF $K_R(S^1 \times D^2, 2)$

Trivalent graphs will be interpreted as in [A2, A3, GH, KL, MV]. Any unlabelled edge is assumed to be colored one. The colors of the three edges incident to a single vertex must form an admissible triple. Given non-negative integers a , b , and c , the triple (a, b, c) is admissible





FIGURE 3. The doubling pairing of two graph basis elements. The bold loop indicates a 0-framed surgery.

if $|a - b| \leq c \leq a + b$ and $a + b + c \equiv 0 \pmod{2}$. We use the notation of [KL]: Δ_n , $\theta(a, b, c)$, $\text{Tet} \begin{bmatrix} a & b & e \\ c & d & f \end{bmatrix}$, and $\lambda_c^{a,b}$.

We use the graph basis defined in [A2, A3]. Given a pair of non-negative integers (i, ε) such that $\varepsilon = i + 1$ or $\varepsilon = i - 1$, let



Let R denote the ring $\mathbb{Z}[A, A^{-1}]$ localized by inverting $A^k - 1$ for all natural numbers k , and let $K_R(M, m)$ denote the Kauffman bracket skein module of M relative to m points with coefficients in R . According to [P, Theorem 2.3], we have that $K_R(M, m) = K(M, m) \otimes R$, so we can essentially view $K(M, m)$ as a subset of $K_R(M, m)$. We make this distinction because when computing a finite list of generators for the even and odd Kauffman bracket ideals, we pass through $K_R(S^1 \times D^2, 2)$ when using the doubling pairing defined in [A2, §2.3]. However, each of the generators we obtain is in fact an element of $K(S^1 \times D^2, 2)$.

Recall, according to [HP], $K_R(S^1 \times S^2)/\text{torsion}$ is isomorphic to R . Let $\psi : K_R(S^1 \times S^2) \rightarrow R$ be the epimorphism that sends the empty link to $1 \in R$. The doubling pairing is defined to be the symmetric pairing $\langle \cdot, \cdot \rangle_D : K_R(S^1 \times D^2, 2) \times K_R(S^1 \times D^2, 2) \rightarrow R$ obtained by gluing two solid tori containing skein elements together via a certain orientation-reversing homeomorphism to obtain a skein element in $S^1 \times S^2$, and evaluating this skein element under ψ . Figure 3 illustrates the doubling pairing of two graph basis elements. The thick dark colored loop indicates where a 0-framed surgery is to be performed, converting S^3 to $S^1 \times S^2$. According to [A2, Theorem 2.4], the graph basis is orthogonal with respect to the doubling pairing.

5. APPLICATIONS TO GENUS-1 TANGLE EMBEDDING

Let \mathcal{G} be a genus-1 tangle. We follow the same basic procedure as in [A2, A3] to obtain finite lists of generators for $I_{\mathcal{G}}^{\text{even}}$ and $I_{\mathcal{G}}^{\text{odd}}$. First, we write \mathcal{G} as a linear combination of graph basis elements $\mathcal{G} = \sum c_{i,\varepsilon} g_{i,\varepsilon}$. Since the graph basis is orthogonal, we have that $c_{i,\varepsilon} = \langle \mathcal{G}, g_{i,\varepsilon} \rangle_D / \langle g_{i,\varepsilon}, g_{i,\varepsilon} \rangle_D$ and only finitely many $c_{i,\varepsilon}$ are non-zero.

We then use this linear combination to compute the relative Hopf pairing of \mathcal{G} with the even (respectively, odd) basis to obtain a generating set for $I_{\mathcal{G}}^{\text{even}}$ (respectively, $I_{\mathcal{G}}^{\text{odd}}$). The following results allow us to compute the relative Hopf pairing of the graph basis with the even and odd bases. The proof of Lemma 5.1 below is similar to that of [A2, Lemma 4.1].

Lemma 5.1.

$$\left. \begin{array}{c} | \\ \text{---} \circ \text{---} \\ | \end{array} \right|_i S_n = \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2) \left| \begin{array}{c} | \\ \text{---} \circ \text{---} \\ | \end{array} \right|_i$$

If $n = 0$, we interpret $\prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2)$ as 1.

Notice that $\prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2)$ is zero if $n > i$.

Proposition 5.2. (i) $\langle g_{i,\varepsilon}, x_n^{\text{even}} \rangle = \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2)$.

$$(ii) \langle g_{i,\varepsilon}, y_n^{\text{even}} \rangle = \phi_i (\lambda_\varepsilon^{i-1})^{-2} \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2).$$

$$(iii) \langle g_{i,\varepsilon}, x_n^{\text{odd}} \rangle = \phi_i \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2).$$

$$(iv) \langle g_{i,\varepsilon}, y_n^{\text{odd}} \rangle = (\lambda_\varepsilon^{i-1})^{-2} \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2).$$

Each of these is zero if $n > i$.

Proof. (i) We have from Lemma 5.1 that

$$\langle g_{i,\varepsilon}, x_n^{\text{even}} \rangle = S_n \left(\begin{array}{c} \text{---} \circ \text{---} \\ | \end{array} \right)_i = \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2) \left(\begin{array}{c} \text{---} \circ \text{---} \\ | \end{array} \right)_i = \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2).$$

(ii) Using Lemma 5.1, [A2, equations 2.2, 2.5] and $\lambda_k^{i,j} = \lambda_k^{j,i}$,

$$\begin{aligned}
\langle g_{i,\varepsilon}, y_n^{\text{even}} \rangle &= \langle \text{diagram with } zS_n \rangle = \phi_i \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2) \langle \text{diagram with } i, \varepsilon \rangle \\
&= \phi_i (\lambda_\varepsilon^{i,1})^{-1} (\lambda_\varepsilon^{1,i})^{-1} \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2) \langle \text{diagram with } i, \varepsilon \rangle \\
&= \phi_i (\lambda_\varepsilon^{i,1})^{-2} \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2).
\end{aligned}$$

(iii)

$$\begin{aligned}
\langle g_{i,\varepsilon}, x_n^{\text{odd}} \rangle &= \langle \text{diagram with } zS_n \rangle = \phi_i \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2) \langle \text{diagram with } i, \varepsilon \rangle \\
&= \phi_i \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2).
\end{aligned}$$

(iv)

$$\begin{aligned}
\langle g_{i,\varepsilon}, y_n^{\text{odd}} \rangle &= \langle \text{diagram with } S_n \rangle = \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2) \langle \text{diagram with } i, \varepsilon \rangle \\
&= (\lambda_\varepsilon^{i,1})^{-1} (\lambda_\varepsilon^{1,i})^{-1} \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2) \langle \text{diagram with } i, \varepsilon \rangle \\
&= (\lambda_\varepsilon^{i,1})^{-2} \theta(1, i, \varepsilon) \prod_{k=0}^{n-1} (\phi_i^2 - \phi_k^2).
\end{aligned}$$

□

Recall, the Kauffman bracket polynomial of any even closure L of \mathcal{G} can be written as $\langle L \rangle = \langle \mathcal{G}, \mathcal{G}' \rangle$ where $\mathcal{G}' \in K^{\text{even}}(S^1 \times D^2, 2)$. So, $\langle \mathcal{G}, x_n^{\text{even}} \rangle / \delta$ and $\langle \mathcal{G}, y_n^{\text{even}} \rangle / \delta$ form a generating set for $I_{\mathcal{G}}^{\text{even}}$. Proposition 5.2 implies that only finitely many of these will be non-zero. Similarly, $\langle \mathcal{G}, x_n^{\text{odd}} \rangle / \delta$ and $\langle \mathcal{G}, y_n^{\text{odd}} \rangle / \delta$ form a finite generating set for $I_{\mathcal{G}}^{\text{odd}}$.

6. EXAMPLES

We compute the even and odd Kauffman bracket ideals for three tangles \mathcal{A} , \mathcal{D} , and \mathcal{H} . In each of these computations, our first step is to compute the doubling pairing of the tangle in question with the graph basis. We leave out the full computation for the sake of brevity, but we follow the same procedure as in [A2, Appendix A]. We then write each tangle as a linear combination of graph basis elements. It turns out that due to admissibility conditions, \mathcal{A} , \mathcal{D} , and \mathcal{H} may all be written as $c_{0,1}g_{0,1} + c_{2,1}g_{2,1} + c_{2,3}g_{2,3}$ for some coefficients $c_{i,\varepsilon} \in R$. Thus we have using Proposition 5.2:

Lemma 6.1. *If $\mathcal{G} = \mathcal{A}$, \mathcal{D} , or \mathcal{H} , $I_{\mathcal{G}}^{\text{even}}$ is generated by $\langle \mathcal{G}, x_i^{\text{even}} \rangle / \delta$ and $\langle \mathcal{G}, y_i^{\text{even}} \rangle / \delta$ where $0 \leq i \leq 2$. Similarly, $I_{\mathcal{G}}^{\text{odd}}$ is generated by $\langle \mathcal{G}, x_i^{\text{odd}} \rangle / \delta$ and $\langle \mathcal{G}, y_i^{\text{odd}} \rangle / \delta$ where $0 \leq i \leq 2$.*

For $\mathcal{G} = \mathcal{A}$, \mathcal{D} or \mathcal{H} , we followed the same procedure as in [A2, A3], to find $I_{\mathcal{A}}^{\text{even}}$ and $I_{\mathcal{A}}^{\text{odd}}$, using Proposition 5.2, Lemma 6.1, and Mathematica. One can verify directly that the claimed ideals are indeed non-trivial using the computations in §7.

6.1. Krebs' Tangle \mathcal{A} . We consider the genus-1 tangle given by Krebs [K] pictured in Figure 1. We have that

$$\langle \mathcal{A}, g_{i,\varepsilon} \rangle_D = \text{Diagram} = \sum_{j,k,l} \frac{\lambda_i^{1-1}(\lambda_j^{1-1})^{-1}(\lambda_k^{1-1})^{-1}(\lambda_l^{1-1})^{-1} \Delta_j \Delta_k \Delta_l \text{Tet} \begin{bmatrix} 1 & i & \varepsilon \\ 1 & j & 1 \end{bmatrix} \text{Tet} \begin{bmatrix} l & 1 & j \\ 1 & k & 1 \end{bmatrix} \text{Tet} \begin{bmatrix} 1 & \varepsilon & 1 \\ k & l & j \end{bmatrix}}{\theta(1,1,i)\theta(1,1,j)\theta(1,1,k)\theta(1,1,l)\theta(1,j,\varepsilon)\theta(\varepsilon,k,1)\theta(l,k,j)}$$

where the sum is over all j , k , and l such that the following triples are admissible: $(1, 1, i)$, $(1, 1, j)$, $(1, j, \varepsilon)$, $(1, 1, k)$, $(\varepsilon, k, 1)$, $(1, 1, l)$, and (l, k, j) . Admissibility conditions imply that 0 and 2 are the only possible admissible values for j , k , and l . Note that, if $i \neq 0, 2$, there are no such j, k, l , and the given sum is over an empty index set. Thus, the value of the sum is zero. So $\langle \mathcal{A}, g_{i,\varepsilon} \rangle_D = 0$ unless $i = 0$ or $i = 2$.

The coefficients for \mathcal{A} as a linear combination of the graph basis are $c_{0,1} = \frac{-1-A^8+A^{12}}{1+A^4}$, $c_{2,1} = \frac{-1+A^4+A^{12}}{A^6+A^{10}+A^{14}}$, and $c_{2,3} = 1$.

After further computation, we obtain the generating sets given in the following result.

Proposition 6.2. *The even Kauffman bracket ideal $I_{\mathcal{A}}^{\text{even}}$ of Krebs' tangle \mathcal{A} is trivial. The odd Kauffman bracket ideal of \mathcal{A} is $I_{\mathcal{A}}^{\text{odd}} = \langle 9, 4 + A^4 \rangle$ which is non-trivial.*

6.2. A small tangle, \mathcal{D} . We now consider the genus-1 tangle \mathcal{D} pictured in Figure 4. In contrast to Krebs' example, \mathcal{D} has a non-trivial even Kauffman bracket ideal and a trivial odd Kauffman bracket ideal.




FIGURE 4. A genus-1 tangle, denoted by \mathcal{D} .

We have that

$$\langle \mathcal{D}, g_{i,\varepsilon} \rangle_D = \text{diagram} \quad \varepsilon =$$

$$\sum_j \frac{\lambda_i^{1-1} (\lambda_j^{1-1})^{-3} \Delta_j \text{Tet} \begin{bmatrix} 1 & 1 & j \\ 1 & \varepsilon & i \end{bmatrix} \text{Tet} \begin{bmatrix} 1 & i & \varepsilon \\ 1 & j & 1 \end{bmatrix}}{\theta(1,1,i)\theta(1,1,j)\theta(1,\varepsilon,j)}$$



where the sum is over all integers j , such that the following are admissible triples: $(1, 1, i)$, $(1, 1, j)$, and $(1, \varepsilon, j)$. Admissibility conditions imply that 0 and 2 are the only possible admissible values for j , and $\langle \mathcal{D}, g_{i,\varepsilon} \rangle_D = 0$ unless $i = 0$ or $i = 2$.

The coefficients for \mathcal{D} as a linear combination of the graph basis are $c_{0,1} = \frac{1-A^4-A^{12}}{A^2+A^6}$, $c_{2,1} = \frac{1+A^8-A^{12}}{A^8+A^{12}+A^{16}}$, and $c_{2,3} = A^2$.

We obtain the following generating sets after further computation.

Proposition 6.3. *The even Kauffman bracket ideal of \mathcal{D} is $I_{\mathcal{D}}^{\text{even}} = \langle -9, -2 + A^4 \rangle$ which is non-trivial. The odd Kauffman bracket ideal $I_{\mathcal{D}}^{\text{odd}}$ of \mathcal{D} is trivial.*

Indeed, one can see that the odd closure shown in Figure 5 is the unknot, so $I_{\mathcal{D}}^{\text{odd}}$ must be trivial.

FIGURE 5. A trivial odd closure of the tangle \mathcal{D} .FIGURE 6. A genus-1 tangle, denoted by \mathcal{H} .

6.3. A particularly interesting tangle, \mathcal{H} . We consider the genus-1 tangle \mathcal{H} pictured in Figure 6. We have that

$$\langle \mathcal{H}, g_{i,\varepsilon} \rangle_D = \text{Diagram} =$$

$$\sum_{j \ k \ l} \frac{\lambda_i^{-1} (\lambda_j^{-1})^{-3} (\lambda_k^{-1})^{-3} \lambda_l^{-1} \Delta_j \Delta_k \Delta_l \text{Tet} \begin{bmatrix} 1 & i & \varepsilon \\ 1 & j & 1 \end{bmatrix} \text{Tet} \begin{bmatrix} \varepsilon & i & 1 \\ 1 & k & 1 \end{bmatrix} \text{Tet} \begin{bmatrix} 1 & 1 & l \\ 1 & \varepsilon & j \end{bmatrix} \text{Tet} \begin{bmatrix} 1 & k & \varepsilon \\ 1 & l & 1 \end{bmatrix}}{\theta(1,1,i)\theta(1,1,j)\theta(1,1,k)\theta(1,1,l)\theta(1,j,\varepsilon)\theta(1,k,\varepsilon)\theta(1,l,\varepsilon)}$$

where the sum is over all j , k , and l such that the following triples are admissible: $(1, 1, i)$, $(1, 1, j)$, $(1, j, \varepsilon)$, $(1, 1, k)$, $(1, k, \varepsilon)$, $(1, 1, l)$, and $(1, l, \varepsilon)$. Admissibility conditions imply that 0 and 2 are the only possible admissible values for j , k , and l . So, $\langle \mathcal{H}, g_{i,\varepsilon} \rangle_D = 0$ unless $i = 0$ or $i = 2$.

The coefficients for \mathcal{H} as a linear combination of the graph basis are

$$c_{0,1} = \frac{-1+2A^4-3A^8+2A^{12}-3A^{16}+2A^{20}-A^{24}+A^{28}}{A^{12}+A^{16}},$$

$$c_{2,1} = \frac{-1+A^4-2A^8+3A^{12}-2A^{16}+3A^{20}-2A^{24}+A^{28}}{A^{18}+A^{22}+A^{26}}, \text{ and } c_{2,3} = A^4.$$

The following generating sets are obtained after further computation.

Proposition 6.4. *The even Kauffman bracket ideal of \mathcal{H} is $I_{\mathcal{H}}^{\text{even}} = \langle 5, 1 + A^4 \rangle$ which is non-trivial. The odd Kauffman bracket ideal of \mathcal{H} is $I_{\mathcal{H}}^{\text{odd}} = \langle 9, 4 + A^4 \rangle$ which is also non-trivial.*

These corollaries follow immediately.

Corollary 6.5. *The Kauffman bracket ideal $I_{\mathcal{H}}$ of the genus-1 tangle \mathcal{H} is trivial.*

Corollary 6.6. *The genus-1 tangle \mathcal{H} does not embed in the unknot.*

Although \mathcal{H} is not obstructed from embedding in the unknot by the ordinary Kauffman bracket ideal, the even and odd Kauffman bracket ideals, working together, do provide an obstruction.

7. RELATION TO DETERMINANTS

Let ω denote $e^{\frac{\pi i}{4}}$, and $\Omega : \mathbb{Z}[A, A^{-1}] \rightarrow \mathbb{Z}[\omega]$ be the ring epimorphism sending A to ω . Recall that the determinant $\det(L)$ of a link L is the absolute value of the determinant of a Seifert matrix for L symmetrized or, equivalently, the order of the first homology of the double branched cover of S^3 along L . If this homology group is infinite, the order is interpreted to be zero. One has that [K, Prop. 1], $\det(L) = \omega^j \Omega(\langle L \rangle')$ for an integer j , chosen so that $\omega^j \Omega(\langle L \rangle')$ is a non-negative integer.

Proposition 7.1. *Let L be a closure of a genus-1 tangle \mathcal{G} . Then $\det(L) \in \Omega(I_{\mathcal{G}}) \cap \mathbb{Z}$. If L is an even closure, $\det(L) \in \Omega(I_{\mathcal{G}}^{\text{even}}) \cap \mathbb{Z}$. If L is an odd closure, $\det(L) \in \Omega(I_{\mathcal{G}}^{\text{odd}}) \cap \mathbb{Z}$.*

Let $\langle n \rangle_{\mathbb{Z}}$ denote the \mathbb{Z} -ideal generated by n . For the ideals computed in the examples, noting that $\Omega(A^4) = -1$, we have:

$$\begin{aligned} \Omega(I_{\mathcal{A}}^{\text{odd}}) \cap \mathbb{Z} &= \Omega(\langle 9, 4 + A^4 \rangle) \cap \mathbb{Z} = \Omega(\langle 3 \rangle) \cap \mathbb{Z} = \langle 3 \rangle_{\mathbb{Z}}. \\ \Omega(I_{\mathcal{H}}^{\text{even}}) \cap \mathbb{Z} &= \Omega(\langle 9, 4 + A^4 \rangle) \cap \mathbb{Z} = \Omega(\langle 3 \rangle) \cap \mathbb{Z} = \langle 3 \rangle_{\mathbb{Z}}. \\ \Omega(I_{\mathcal{D}}^{\text{even}}) \cap \mathbb{Z} &= \Omega(\langle 9, -2 + A^4 \rangle) \cap \mathbb{Z} = \Omega(\langle 3 \rangle) \cap \mathbb{Z} = \langle 3 \rangle_{\mathbb{Z}}. \\ \Omega(I_{\mathcal{H}}^{\text{odd}}) \cap \mathbb{Z} &= \Omega(\langle 5, 1 + A^4 \rangle) \cap \mathbb{Z} = \Omega(\langle 5 \rangle) \cap \mathbb{Z} = \langle 5 \rangle_{\mathbb{Z}}. \end{aligned}$$

The first equation above gives a second proof of a result in [A, Theorem 1.3] which is the first sentence in the Proposition 7.2 below. Proposition 7.2 follows from the above equations and Proposition 7.1.

Proposition 7.2. *Let L be an odd closure of \mathcal{A} , then $\det(L) \equiv 0 \pmod{3}$. Let L be an even closure of \mathcal{D} , then $\det(L) \equiv 0 \pmod{3}$. If L is an odd closure of \mathcal{H} , $\det(L) \equiv 0 \pmod{5}$. If L is an even closure of \mathcal{H} , $\det(L) \equiv 0 \pmod{3}$.*

Note for the tangle \mathcal{F} shown in [A2] to have $I_{\mathcal{F}} = \langle 11, 4 - A^4 \rangle$, one has $\Omega(I_{\mathcal{F}}) \cap Z = \langle 1 \rangle_{\mathbb{Z}}$.

REFERENCES

- [A] S.M. Abernathy. On Krebs's tangle. *Topology Appl.* **160** (2013) 1379–1383.
- [A2] S.M. Abernathy. The Kauffman bracket ideal for genus-1 tangles, *J. Knot Theory Ramifications* **24** (2015), no. 2, 1550007.
- [A3] S.M. Abernathy. Obstructions to Embedding Genus-1 Tangles in Links, Thesis (Ph.D.), Louisiana State University (2014). <http://etd.lsu.edu/docs/available/etd-07042014-141943/>
- [BHMV] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel. Three manifold invariants derived from the Kauffman bracket, *Topology* **31** (1992) 685–699.
- [GH] P.M. Gilmer and J. Harris. On the Kauffman bracket skein module of the quaternionic manifold, *J. Knot Theory Ramifications* **16** (2007) 103–125.
- [Hab] K. Habiro. On the colored Jones polynomials of some simple links, *Sūrikaiseikikenkyūsho Kōkyūroku* **1172** (2000) 34–43.
- [HP] J. Hoste, J.H. Przytycki. The Kauffman bracket skein module of $S^1 \times S^2$, *Math. Zeit.* **220** (1995), 65–73
- [KL] L.H. Kauffman and S.L. Lins, *Temperley-Lieb Recoupling Theory and Invariants of 3-manifolds* (Princeton University Press, Princeton, NJ, 1994).
- [K] D.A. Krebs, An obstruction to embedding 4-tangles in links, *J. Knot Theory and its Ramifications* **8** (1999) 321–352.
- [MV] G. Masbaum and P. Vogel. 3-valent graphs and the Kauffman bracket. *Pacific J. Math.* **164** (1994) 361–381.
- [P] J.H. Przytycki, Fundamentals of Kauffman bracket skein modules, *Kobe J. Math* (1999) 45–66.
- [PSW] J.H. Przytycki, D.S. Silver and S.G. Williams, 3-manifolds, tangles, and persistent invariants, *Math. Proc. Camb. Phil. Soc.* **139** (2005) 291–306.

DEPARTMENT OF MATHEMATICS, ANGELO STATE UNIVERSITY, ASU STATION #10900, SAN ANGELO, TX 76909, USA

E-mail address: susan.abernathy@angelo.edu

URL: <http://www.angelo.edu/faculty/sabernathy/>

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA

E-mail address: gilmer@math.lsu.edu

URL: www.math.lsu.edu/~gilmer/